

CME Article

Hiv and Hepatitis C Co-Infection

JAMES T. DWYER, DO, AND SINDY M. PAUL, MD, MPH

Infections with HIV and hepatitis C (HCV) are individually severe diseases that call for complex medical managements. HIV–HCV co-infection complicates the medical management of both diseases. This paper provides an overview of HIV–HCV co-infection with a brief review of the natural history of hepatitis C as it applies to the co-infected patient and recommendations for the diagnosis and medical management of the co-infected patient.

LEARNING OBJECTIVES

- I. To review the epidemiology of HIV–HCV co-infection.
- II. To recognize the need to assess HIV infected patients for HCV co-infection.
- III. To describe a treatment regimen for HIV–HCV co-infected patients.

Hepatitis C (HCV) is one of the most frequent infections related to chronic liver disease and currently the most common blood-borne pathogen in the United States. Recent Centers for Disease Control and Prevention (CDC) data estimates 3.9 million Americans have been infected by hepatitis C to date, and approximately 2.7 million of this group have chronic infection with an estimated annual number of chronic liver disease deaths related to hepatitis C ranging from 8,000–10,000. Current CDC disease burden data estimates that 25,000 newly acquired infections have been reported as of 2001, and 4000 acute clinical cases were reported during the same time period. The 1.8% of the general population have at some time been infected with hepatitis C.¹

The prevalence of HCV co-infection in HIV patients averages 35%. But in HIV clinic populations, where there is a high prevalence of injection drug use (IDU) as a risk factor for acquiring HIV, the prevalence may be as high as 80%–90%. High prevalence rates (> 50%) have also been reported in HIV positive inmates at correctional facilities and among patients treated for hemophilia.² Currently, 150,000–300,000 persons in the United States are infected with both HIV and hepatitis C. This number represents 15%–30% of all HIV-infected persons by some estimates, and 5%–10% of all HCV-infected persons.³

HEPATITIS C DISEASE

Hepatitis C is a single stranded RNA virus of the Flaviviridae family. Currently, there are six known genotypes and more than fifty serotypes of this virus. The different genotypes have different worldwide geographic distribution. In the United States, genotype 1 accounts for 75% or more of all HCV infections and is reported to have the poorest response to therapy of all the genotypes.⁴

JAMES T. DWYER, DO, is a Fellow in Infectious Diseases at UMDNJ–School of Osteopathic Medicine/Garden State Infectious Disease Associates, and is a Master of Public Health student at UMDNJ, Stratford. SINDY M. PAUL, MD, MPH, is medical director of the Division of AIDS Prevention and Control, New Jersey Department of Health and Senior Services.

DISCLOSURE STATEMENT: James T. Dwyer, DO, and Sindy M. Paul, MD, MPH, have no relationships to disclose. The article includes a discussion of commercial products or services.

Hcv is spread predominantly by contact with infected blood or blood products. Prior to 1992, blood transfusion was a leading cause for HCV transmission. Since that time, blood donations are routinely screened for the presence of anti-HCV. Today, the use of shared, unsterilized needles or syringes by injection-drug users; occupational exposure; and, to a lesser extent, maternal-fetal and sexual transmission have been reported as causes of HCV infection.^{3,5,6,7} When looking at risk factors for co-infection with HIV, the CDC estimates are that from 50%–90% of HIV-infected drug users are co-infected with HCV.⁴ The risk of acquiring HCV through perinatal or sexual exposures is much lower than it would be for HIV and estimates range from 3%–5% risk of transmission.⁵ Sporadic transmission, defined as exposure to the virus from cuts, wounds, needle sticks, or procedures occurs in about 10% of acute hepatitis c and in 30% of chronic hepatitis c cases.⁶

Chronic hepatitis c varies greatly in its course and outcome. There are patients who have no signs and symptoms of liver disease. Liver biopsy usually shows some degree of chronic hepatitis, but the degree of liver injury is usually mild, and the overall prognosis may be good. On the other hand, there are patients with severe, symptomatic hepatitis with detectable HCV RNA in serum, and concurrent elevation of hepatic enzymes ultimately evolving to end-stage liver disease, cirrhosis, or hepatic cell carcinoma (HCC).⁶ Although approximately 20% of persons who are infected with HCV alone clear HCV RNA from their blood after acute infection, HCV clearance occurs in only 5%–10% of HIV infected persons, less frequently in those with lower CD4+ cell counts.³ In the co-infected patient, HCV disease progression is enhanced, and end-stage liver disease and cirrhosis may present without significant signs or symptoms heralding its presence.

HIV–HCV CO-INFECTION

It is important to understand how HIV and HCV interact in the co-infected patient. Hcv has a direct impact on HIV disease. The phenomenon of co-in-

fection results in more cases of end-stage liver disease and cirrhosis than in HCV alone.^{3,7,8,9} HCV CD4+ cell increase may be smaller in the co-infected patient, even with effective anti-HIV therapy. This suggests that HCV may blunt immune recovery. Infection with HIV has been associated with higher HCV RNA viral loads and, in most studies, with more rapid progression of cirrhosis, liver failure, and HCC.³ Hepatotoxicity caused by antiretroviral drugs is always a concern in a patient with liver disease. The co-infected patient displays greater morbidity and mortality from HCV than from HIV disease alone. Although, highly active antiretroviral therapy (HAART) improves patients' overall HIV indicators, no significant change occurs in alanine aminotransferase (ALT) and HCV RNA levels.¹⁰ In the era of HAART, clinicians now must concern themselves with addressing the co-morbidities of patients living with HIV–AIDS and HCV.

DIAGNOSIS

The diagnosis of hepatitis c should be entertained in all HIV positive patients. In order to formulate a care plan, a complete hepatitis profile is advisable for all HIV positive patients. Liver enzyme abnormalities alone are not an accurate predictor of infection with hepatitis c and should not be relied upon as a screening test for HCV or other types of hepatitis. Initially, screening for HCV should include an HCV Enzyme Immunoassay (EIA). If the HCV EIA is negative in a patient at risk for HCV, a PCR for HCV RNA should be considered. A patient with a positive HCV EIA should have an HCV RNA PCR to confirm the presence of viremia.² If viremia is evident, an HCV genotype is warranted. This will provide useful information in planning future treatment options. Because there are no clinical, biochemical, or virologic factors that could reliably identify patients with advanced fibrosis or cirrhosis,⁷ a liver biopsy in all co-infected patients is needed. In addition, if treatment of HCV is being considered, thyroid stimulating hormone (TSH), complete blood count (CBC), prothrombin time (PT), ferritin, alpha-fetoprotein (AFP), anti-nuclear antibody (ANA), and liver function studies (LFTs) should be part of the

initial evaluation.² A psychiatric evaluation should be performed because of the increased incidence of neuropsychiatric events observed in patients with chronic diseases and in those patients being treated with interferon (IFN) for HCV.

TREATMENT

In 2002 the NIH consensus statement recommended that, although there were no specific HCV therapies approved for the co-infected patient, these patients should be considered for treatment.⁴ Currently, there are no therapies that have been approved by the Food and Drug Administration (FDA) for the treatment of the HIV–HCV co-infected patient, although preliminary studies suggest the combination of Pegylated Alpha Interferon (PEG-Intron A)/Ribavirin (Rebtrrol) is an optimal therapy in the co-infected patient.^{3,6} The recommended dose of IFN and Ribavirin are based on body weight. In combination therapy, the recommended doses of PEG-Intron are 1.5 micrograms per kilogram per week, in a single weekly dose. The recommended Rebtrrol dose is 600–800 mg twice a day. Dose reductions are recommended in patients with renal failure.¹¹

PEG-Intron is contraindicated in those patients with autoimmune hepatitis or decompensated liver disease. Additionally, combination therapy is contraindicated in patients with hemoglobinopathies, in women who are pregnant, and in men whose female partner is pregnant. If a man with hepatitis C is being treated and his female partner is pregnant, teratogenic effects to the fetus can be observed secondary to the teratogenic effects of ribavirin.¹¹

The patient should be closely monitored during therapy for adverse reactions, some of which may be life threatening. They include suicidal–homicidal ideation, depression–anxiety, psychosis, and relapsing substance abuse issues, which underscores the importance of a psychiatric evaluation prior to initiation of therapy. The treatment may aggravate thyroid disorders and precipitate hyperglycemia. Pancreatitis and bone marrow suppression has been observed. Hypotension, cardiac

arrhythmia, and myocardial infarction with IFN have been reported. Pulmonary disorders including pneumonia, dyspnea, and interstitial pneumonitis may be induced or aggravated with therapy. Abdominal pain and fatal ulcerative–ischemic colitis may be seen within the first three months of therapy. Exacerbation or development of autoimmune disorders may occur, and various ophthalmologic conditions have been reported. Rarely, hypersensitivity reactions have occurred. Constitutional symptoms of myalgia, headache, nausea, vomiting, fatigue, and other flu-like symptoms are expected findings at the initiation of therapy, but should subside within a few weeks.

Concomitant use of benzodiazepines may result in hepatotoxicity. Methadone has been found to be less effective in those who are receiving IFN therapy. Various nucleoside reverse transcriptase analogues (NRTI) have been found to interact with IFN, although to date, clinically significant antagonism has not been documented.^{3,11}

The initiation of HAART may be limited by either HCV-related liver disease or the hepatotoxicity of the medications in concert with viral liver disease.¹² Overall, researchers have found that there were proportionately more deaths in the HIV–HCV cohort group when compared to a control group of HIV mono-infected patients' survival;⁸ however, the co-infected patient receiving HAART had similar durations of survival when compared to the HIV mono-infected patient. This implies that managing HIV disease plays a major role in overall survival in the co-infected patient and that managing the potential hepatotoxicities of these medications continues to be a major challenge to clinicians.

Histological improvement on repeat liver biopsy was frequently observed in patients receiving interferon therapy, even without the absence of HCV clearance.¹³ A small cohort study in Japan also found that in the co-infected patient, the use of high doses of interferon reduced viral burden of both HIV and HCV.¹⁴ An important finding of the AIDS Clinical Trial Group (ACTG) protocol A5071, is that, while there were more toxicities associated with the use of PEG-Intron, no negative impact was seen on

parameters used for monitoring HIV disease.¹⁵ There are on-going studies to evaluate the safety, efficacy, and pharmacokinetics of combination HCV therapy in conjunction with HAART, but caution should be exercised when using simultaneous treatments. Treatment of HCV and HIV should not be initiated simultaneously.

To date there is no solid consensus to the answer of the question, “Which disease should be treated first, HIV or HCV?” Surely this is a fertile area for future research. Certainly it has become evident that the progression of HCV in the co-infected patient is more rapid than in the mono-infected HCV patient, and mortality is improved with initiation and use of HAART in the co-infected patient.⁸ It is of the utmost importance that in a co-infected patient, both disease entities be serologically and histologically pursued before any treatment regimen is contemplated. The potential toxicities and interactions of the medications used must be anticipated and minimized to assure the most optimal outcome, and patients must be educated accordingly. If there is relative stability in a patient’s HIV disease, (CD4+ \geq 350, HIV RNA \leq 55,000), it may be more prudent to treat HCV first; however, if this is not the case, considering the initiation of HAART is warranted.

LABORATORY WORK

As with all patients, close and frequent follow-up is a necessity. Initially, a complete blood count and LFTs should be evaluated every 2 weeks for the first month of therapy, and then every 1–2 months thereafter. Dose adjustments of IFN and of Ribavirin may need to be made at that time. HIV RNA and CD4+ cell count should be performed at week 12 of therapy, and an HCV RNA PCR should be measured at week 24. If there is undetectable HCV RNA, therapy is continued for an additional 24 weeks. Some clinicians will stop therapy at this time in persons with HCV genotype 2 or 3; this approach has not been evaluated in the co-infected patient.³ Periodic evaluation of electrolytes and of kidney function is warranted. At the end of therapy and six months thereafter an HCV RNA PCR should be completed. If the HCV RNA is undetectable 6 months after stop-

ping therapy, the chance of virologic relapse is low.³ Data on sustained virologic response (SVR) is not available; however, approximately 23% of co-infected patients had undetectable HCV RNA after 12 weeks of combination therapy.³ Few options exist for patients who either do not respond to therapy, or who respond, and later relapse. The use of long-term therapy/continual therapy for those patients who initially responded to therapy may be an option; however, this requires further investigation.⁴

ADDITIONAL CONSIDERATIONS

In addition to the use of antiviral agents, a patient’s immunization status should be assessed. Immunizations for hepatitis A (HAV) and hepatitis B (HBV) should be given in all patients when appropriate. The pneumococcal and yearly influenza vaccine is appropriate for all patients. Counseling related to the use of dietary supplements, especially vitamin A or iron should be addressed, as these substances accumulate in the liver. Abstinence from alcohol intake or other substance abuse must be stressed.

SUMMARY

Understanding of the co-infected patient continues to evolve. Currently the majority of research, diagnosis, and treatment options have been geared to the mono-infected HIV or HCV patient. The dynamics created by the interactions of these two disease entities requires the clinician to understand the complexity of evaluating, diagnosing, and treating the co-infected patient. Future research should be designed to evaluate novel, safe, and efficacious therapies to optimize the outcome in this growing population. *NJM*

REFERENCES

1. Centers for Disease Control and Prevention—National Center for Infectious Diseases. Viral Hepatitis Surveillance. Available at URL: http://www.cdc.gov/ncidod/diseases/hepatitis/resource/dz_burden02.htm
2. “Expert Perspective III: Strategies for the Management of HIV/HCV Coinfection” (2002). Education Initiative in Infectious Disease—Projects in Knowledge.

3. M. Sulkowski et al. "Hepatitis C in the Hiv Infected Person," *Annals of Internal Medicine* 138 (2003): 197–207.
4. National Institutes of Health–Consensus Development Conference Statement. "Management of Hepatitis C: 2002."
5. Centers for Disease Control and Prevention–Divisions of Hiv–Aids Prevention. "Frequently Asked Questions and Answers about Coinfection with Hiv and Hepatitis c Virus". Available at URL: www.cdc.gov/hiv.
6. Chronic Hepatitis c: Current Disease Management. NIDDK–National Digestive Diseases Information Clearinghouse. Available at URL: www.niddk.nih.gov/health/digest/d
7. R. Sterling et al. "The Clinical Spectrum of Hepatitis c Virus in Hiv Coinfection," *Journal of Acquired Immune Deficiency Syndromes* 32 (2003): 30–37.
8. E. Tedaldi et al. "Influence of Co-infection with Hepatitis c Virus on Morbidity and Mortality Due to Human Immunodeficiency Virus Infection in the Era of Highly Active Antiretroviral Therapy," *Clinical Infectious Diseases* 36 (2003): 363–367.
9. C. S. Graham et al. "Influence of Human Immunodeficiency Virus Infection on the Course of Hepatitis c Virus Infection: A Meta-Analysis," *Clinical Infectious Diseases* 33 (2001): 562–569.
10. D. Torre et al. "Evolution of Co-infection with Human Immunodeficiency Virus and Hepatitis c Virus in Patients Treated with Highly Active Antiretroviral Therapy," *Clinical Infectious Diseases* 33 (2001): 1579–1585.
11. *Physicians' Desk Reference*, edition 57 (Montvale, N.J.: Thompson PDR Publishing, 2003): 1742.
12. B. McGovern. "Evaluation and Treatment of the Hiv/Hcv Co-infected Patient: A Case History and Discussion," *Aids Patient Care and STDs* 16, no. 7 (2002): 319–326.
13. V. Dimartino et al. "Hiv Co-infection Does Not Compromise Liver Histological Response to Interferon Therapy in Patients with Chronic Hepatitis c," *AIDS* 16 (2002): 441–445.
14. H. Hanabusa. "Efficacy of Induction Therapy with High-Dose Interferon for Patients with Hemophilia and Human Immunodeficiency Virus–Hepatitis c Co-infection," *Clinical Infectious Diseases* 35, no. 12(2002): 1527–1533.
15. K. Sherman et al. "Hepatitis c Virus Prevalence among Patients Infected with Human Immunodeficiency Virus: A Cross-Sectional Analysis of the U.S. Adult Aids Clinical Trials Group," *Clinical Infectious Diseases* 34 (2002): 831–837.

CME Examinations

INSTRUCTIONS

Type or print your full name and address and your date of birth in the spaces provided on the CME REGISTRATION FORM, then record your answers. Retain a copy of your answers.

Complete the evaluation portion of the CME REGISTRATION FORM. Forms and examinations cannot be processed if the evaluation portion is incomplete. Evaluations of this article in no way affect the scoring of the examinations.

Send the completed form to: *New Jersey Medicine*, Medical Society of New Jersey, Two Princess Road, Lawrenceville, NJ 08648; or fax the completed form to: *New Jersey Medicine*, 609-896-1368.

Examinations will be graded, and you will be advised as to whether you have passed or failed. Unanswered questions are considered to be incorrect. A score of at least 70% is required to pass. Answers to the examination will be sent to you along with your CME certificate.

Failed examinations may be retaken.

Be sure to submit the CME REGISTRATION FORM on or before the deadline. Forms received after the listed deadline will not be processed.

CME ACCREDITATION

This CME article is primarily targeted at physicians and other allied health professionals. There are no specific background requirements. This activity has been planned and implemented in accordance

with the Essential Areas and Policies of the Accreditation Council for Continuing Medical Education (ACCME) through the joint sponsorship of the Academy of Medicine of New Jersey (AMNJ) and the Medical Society of New Jersey (MSNJ). AMNJ is accredited by the ACCME to provide continuing medical education for physicians.

CREDIT DESIGNATION

AMNJ designates this article for a maximum of 1.0 hour in category 1 credit towards the AMA Physician's Recognition Award. Each physician should claim only those hours of credit actually spent.

FULL-DISCLOSURE POLICY

It is the policy of the Academy of Medicine of New Jersey to ensure balance, independence, objectivity, and scientific rigor in all of its educational activities. All authors participating in continuing medical education programs sponsored by the Academy of Medicine of New Jersey are expected to disclose to the audience any real or apparent conflict(s) of interest related to the content of their material. Full disclosure of author relationships will be made in the article.

STATEMENT OF COMPLIANCE

This activity has been planned and implemented in accordance with the Essential Areas and Policies of the Accreditation Council for Continuing Medical Education (ACCME) by the Academy of Medicine of New Jersey.

CME EXAMINATION: DEADLINE SEPTEMBER 30, 2004

“Hiv and Hepatitis C Co-infection”

1. What is the prevalence of HIV–HCV co-infection among HIV infected injection drug users?
 - A. 60%–70%
 - B. 70%–80%
 - C. 80%–90%
 - D. 90%–100%
2. Which of the best describes the impact of HCV on HIV disease?
 - A. Blunt CD4+ cell recovery
 - B. Decreases HIV viral load
 - C. Decreases HIV disease progression
 - D. No influence
3. What initial test should be performed in HIV infected patients to assess their HCV status?
 - A. HCV EIA
 - B. HCV RNA
 - C. Liver biopsy
 - D. Liver function studies
4. Which of the following is a contraindication for pegylated interferon?
 - A. Hepatitis B infection
 - B. HIV infection
 - C. Pregnancy
 - D. Renal disease
5. At what point during therapy should HCV RNA be re-evaluated?
 - A. 12 weeks
 - B. 24 weeks
 - C. 36 weeks
 - D. 48 weeks

ANSWER SHEET

“Hiv and Hepatitis C Co-infection”

Darken the correct answers

1. ☐ A ☐ B ☐ C ☐ D2. ☐ A ☐ B ☐ C ☐ D3. ☐ A ☐ B ☐ C ☐ D4. ☐ A ☐ B ☐ C ☐ D5. ☐ A ☐ B ☐ C ☐ D

Time spent reading this article and completing the learning assessment and evaluation: _____HOURS _____MINUTES

EVALUATION FORM

(This must be completed for this examination to be scored.)

“Hiv and Hepatitis C Co-infection”

Check the appropriate answer below

YES NO

The objectives were useful in determining if this activity would be a worthwhile educational activity for me.

The objectives accurately described the content of and potential learning from the article.

This article will help to modify my practice performance.

The quiz questions were at an appropriate level for assessing my learning.

DEADLINE FOR MAILING: For credit to be received, the envelope must be postmarked no later than **September 30, 2004**.

RETAIN A COPY OF YOUR ANSWERS and compare them with the correct answers, which will be sent with your certificate.

REGISTRATION FORM

(please print or type)

LAST NAME

FIRST NAME

DEGREE

MAILING ADDRESS

CITY

STATE

ZIP CODE

DATE OF BIRTH (USED FOR TRACKING CREDITS ONLY)

PHONE NUMBER

FAX NUMBER

E-MAIL

Send completed form to:

New Jersey Medicine, Medical Society of New Jersey, 2 Princess Road, Lawrenceville, New Jersey 08648,
FAX: 609-896-1368